

# First observation and measurement of the resonant structure of the $\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^- \pi^+ \pi^-$ decay mode

*P. Azzurri<sup>1</sup>, P. Barria<sup>2</sup>, M. A. Ciocci<sup>2</sup>, S. Donati<sup>3</sup> and E. Vataga<sup>1</sup>, for the CDF Collaboration.*

<sup>1</sup>Scuola Normale Superiore, Piazza dei Cavalieri 7, 56126 Pisa, Italy

<sup>2</sup>Università di Siena, Dipartimento di Fisica, Via Roma 56, 53100 Siena, Italy

<sup>3</sup>Università di Pisa, Dipartimento di Fisica, Largo Bruno Pontecorvo 3, 56127 Pisa, Italy

**DOI:** will be assigned

We present the first observation of the  $\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^- \pi^+ \pi^-$  decay using data from an integrated luminosity of approximately  $2.4 \text{ fb}^{-1}$  of  $p\bar{p}$  collisions at  $\sqrt{s}=1.96 \text{ TeV}$ , collected with the CDF II detector at the Fermilab Tevatron. We also present the first observation of the resonant decays  $\Lambda_b^0 \rightarrow \Sigma_c(2455)^0 \pi^+ \pi^- \rightarrow \Lambda_c^+ \pi^- \pi^+ \pi^-$ ,  $\Lambda_b^0 \rightarrow \Sigma_c(2455)^{++} \pi^- \pi^- \rightarrow \Lambda_c^+ \pi^- \pi^+ \pi^-$ ,  $\Lambda_b^0 \rightarrow \Lambda_c(2595)^+ \pi^- \rightarrow \Lambda_c^+ \pi^- \pi^+ \pi^-$  and  $\Lambda_b^0 \rightarrow \Lambda_c(2625)^+ \pi^- \rightarrow \Lambda_c^+ \pi^- \pi^+ \pi^-$ , and measure their relative branching ratios.

## 1 Introduction

Presented here is the observation of the  $\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^- \pi^+ \pi^-$  decay and resonant structure in analogy to the decay structure observed in the  $\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^+ \pi^- \mu^- \bar{\nu}_\mu$  channel [1]. All new measurements of the  $\Lambda_b^0$  branching ratios can be compared to theoretical predictions in the heavy quark effective theory (HQEF) approximation [2].

This measurement is based on data from an integrated luminosity of approximately  $2.4 \text{ fb}^{-1}$  of  $p\bar{p}$  collisions at  $\sqrt{s}=1.96 \text{ TeV}$ , collected with the CDF II detector [3], using two-track impact parameter triggers. Unless stated otherwise, branching fractions, fragmentation functions, and lifetimes used in the analysis are obtained from the Particle Data Group world averages [4].

## 2 Event selection and signal yields

The event reconstruction and selection has been optimized in order to maximize the statistical significance of the total number of  $\Lambda_b^0$  decays observed on the data. The  $\Lambda_c^+$  candidates are reconstructed in the  $\Lambda_c^+ \rightarrow pK^- \pi^+$  channel requiring a vertex  $\chi^2$  probability in excess of  $10^{-4}$ , a transverse decay length in excess of  $200 \mu\text{m}$ ,  $p_T(p) > p_T(\pi^+)$ ,  $p_T(\Lambda_c^+) > 4 \text{ GeV}/c$  and the  $\Lambda_c^+$  invariant mass in the  $2.24\text{--}2.33 \text{ GeV}/c^2$  mass range.

The  $\Lambda_b^0$  candidates are reconstructed by further adding to the  $\Lambda_c^+$  candidates three pion candidate tracks, with  $\eta\phi$ -opening  $\Delta R(3\pi)$  smaller than 1.2. The  $\Lambda_b^0$  candidate is required to have a vertex  $\chi^2$  probability in excess of  $10^{-4}$ , a transverse decay length in excess of  $200 \mu\text{m}$  and a significance in excess of 16, an impact parameter smaller than  $70 \mu\text{m}$ , and a transverse momentum in excess of  $9 \text{ GeV}/c$ .

The resulting distribution of the invariant mass difference  $m(\Lambda_c^+ \pi^- \pi^+ \pi^-) - m(\Lambda_c^+)$  with the  $\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^- \pi^+ \pi^-$  signal peak, is shown in Figure 1. A total signal yield of  $848 \pm 93$  candidates is evaluated with an unbinned likelihood fit using a Gaussian distribution for the signal, an exponential distribution for the background, and Monte Carlo templates for  $B^0$  and  $B_s^0$  backgrounds. In the following  $\Lambda_b^0$  candidates have been selected within  $48 \text{ MeV}/c^2$  of the mass peak.




Figure 1: The reconstructed invariant mass difference  $m(\Lambda_c^+ \pi^- \pi^+ \pi^-) - m(\Lambda_c^+)$ , after applying optimized cuts, showing the total  $\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^- \pi^+ \pi^-$  signal yield.




Figure 2: The reconstructed invariant mass difference  $m(\Lambda_c^+ \pi^- \pi^+) - m(\Lambda_c^+)$  within the  $\Lambda_b^0$  mass window, showing the  $\Lambda_b^0 \rightarrow \Lambda_c(2595)^+ \pi^- \rightarrow \Lambda_c^+ \pi^- \pi^+ \pi^-$  and  $\Lambda_b^0 \rightarrow \Lambda_c(2625)^+ \pi^- \rightarrow \Lambda_c^+ \pi^- \pi^+ \pi^-$  signal yields.

The mass difference  $\Delta m^{-+} = m(\Lambda_c^+ \pi^- \pi^+) - m(\Lambda_c^+)$  for selected  $\Lambda_b^0$  candidates is shown in Figure 2, with the two peaks from  $\Lambda_c(2595)^+$  and  $\Lambda_c(2625)^+$  decays. A fit performed with two signal peaks and a linear background yields  $46.6 \pm 9.7$   $\Lambda_b^0 \rightarrow \Lambda_c(2595)^+ \pi^-$  candidates and  $114 \pm 13$   $\Lambda_b^0 \rightarrow \Lambda_c(2625)^+ \pi^-$  candidates.

Finally the mass differences  $m(\Lambda_c^+ \pi^+) - m(\Lambda_c^+)$  and  $m(\Lambda_c^+ \pi^-) - m(\Lambda_c^+)$  are shown in Figure 2, for selected  $\Lambda_b^0$  candidates, after removing  $\Lambda_c(2595)^+$  and  $\Lambda_c(2625)^+$  decays with the  $\Delta m^{-+} > 360 \text{ MeV}/c^2$  requirement. Separate fits of the two signal contributions yield  $41.5 \pm 9.3$   $\Lambda_b^0 \rightarrow \Sigma_c(2455)^0 \pi^+ \pi^-$  candidates and  $81 \pm 15$   $\Lambda_b^0 \rightarrow \Sigma_c(2455)^{++} \pi^+ \pi^+$  candidates.

### 3 Results

Results are expressed in terms of relative branching fractions between the above resonant decay modes, correcting for the relative channel efficiencies with Monte Carlo simulations. Several sources of systematic effects have been considered, and the dominant uncertainties come from the  $\Lambda_b^0$  and  $\Lambda_c^+$  polarization uncertainty, and on the unknown fraction of non-resonant decays.




Figure 3: The invariant mass difference  $m(\Lambda_c^+ \pi^-) - m(\Lambda_c^+)$  (left) and  $m(\Lambda_c^+ \pi^+) - m(\Lambda_c^+)$  (right) for selected  $\Lambda_b^0$  candidates, after removing events with  $\Lambda_c(2595)^+$  and  $\Lambda_c(2625)^+$  decays, and showing respectively the presence of  $\Lambda_b^0 \rightarrow \Sigma_c(2455)^0 \pi^+ \pi^-$  and  $\Lambda_b^0 \rightarrow \Sigma_c(2455)^{++} \pi^- \pi^-$  signals.

In summary the measured relative branching fractions are the following

$$\begin{aligned}
 \frac{\mathcal{B}(\Lambda_b^0 \rightarrow \Lambda_c(2595)^+ \pi^- \rightarrow \Lambda_c^+ \pi^- \pi^+ \pi^-)}{\mathcal{B}(\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^- \pi^+ \pi^-)} &= (2.5 \pm 0.6(\text{stat}) \pm 0.5(\text{syst})) \times 10^{-2} \\
 \frac{\mathcal{B}(\Lambda_b^0 \rightarrow \Lambda_c(2625)^+ \pi^- \rightarrow \Lambda_c^+ \pi^- \pi^+ \pi^-)}{\mathcal{B}(\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^- \pi^+ \pi^-)} &= (6.2 \pm 1.0(\text{stat})^{+1.0}_{-0.9}(\text{syst})) \times 10^{-2} \\
 \frac{\mathcal{B}(\Lambda_b^0 \rightarrow \Sigma_c(2455)^{++} \pi^- \pi^- \rightarrow \Lambda_c^+ \pi^- \pi^+ \pi^-)}{\mathcal{B}(\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^- \pi^+ \pi^-)} &= (5.2 \pm 1.1(\text{stat}) \pm 0.8(\text{syst})) \times 10^{-2} \\
 \frac{\mathcal{B}(\Lambda_b^0 \rightarrow \Sigma_c(2455)^0 \pi^+ \pi^- \rightarrow \Lambda_c^+ \pi^- \pi^+ \pi^-)}{\mathcal{B}(\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^- \pi^+ \pi^-)} &= (8.9 \pm 2.1(\text{stat})^{+1.2}_{-1.0}(\text{syst})) \times 10^{-2} \\
 \frac{\mathcal{B}(\Lambda_b^0 \rightarrow \Lambda_c(2595)^+ \pi^- \rightarrow \Lambda_c^+ \pi^- \pi^+ \pi^-)}{\mathcal{B}(\Lambda_b^0 \rightarrow \Lambda_c(2625)^+ \pi^- \rightarrow \Lambda_c^+ \pi^- \pi^+ \pi^-)} &= (40.3 \pm 9.8(\text{stat})^{+2.3}_{-1.8}(\text{syst})) \cdot 10^{-2} \\
 \frac{\mathcal{B}(\Lambda_b^0 \rightarrow \Sigma_c(2455)^{++} \pi^- \pi^- \rightarrow \Lambda_c^+ \pi^- \pi^+ \pi^-)}{\mathcal{B}(\Lambda_b^0 \rightarrow \Sigma_c(2455)^0 \pi^+ \pi^- \rightarrow \Lambda_c^+ \pi^- \pi^+ \pi^-)} &= (58.1 \pm 16.9(\text{stat})^{+6.3}_{-9.1}(\text{syst})) \cdot 10^{-2} \\
 \frac{\mathcal{B}(\Lambda_b^0 \rightarrow \Lambda_c(2625)^+ \pi^- \rightarrow \Lambda_c^+ \pi^- \pi^+ \pi^-)}{\mathcal{B}(\Lambda_b^0 \rightarrow \Sigma_c(2455)^{++} \pi^- \pi^- \rightarrow \Lambda_c^+ \pi^- \pi^+ \pi^-)} &= 1.20 \pm 0.26(\text{stat})^{+0.05}_{-0.09}(\text{syst})
 \end{aligned}$$

where the first error is statistical and the second is from systematic uncertainties.

## References

- [1] T. Aaltonen *et al.* (CDF Collaboration), Phys. Rev. **D79** 032001 (2009).
- [2] A. V. Manohar and M. B. Wise, Cambr. Monogr. Part. Phys. Nucl. Phys. Cosmol. 10, 1 (2000); S. Godfrey and N. Isgur, Phys. Rev. **D32**, 189 (1985); N. Isgur, D. Scora, B. Grinstein and M. B. Wise Phys. Rev. **D39**, 799 (1989); A. K. Liebovich, I. W. Stewart, Phys. Rev. **D57**, 5620 (1998); A. K. Liebovich, Z. Ligeti, I. W. Stewart, M. Wise, Phys. Lett **B586**, 337 (2004).
- [3] D. Acosta *et al.* (CDF Collaboration), Phys. Rev. **D71** 032001 (2005).
- [4] C. Amsler *et al.* (Particle Data Group), Phys. Lett. **B667** 1 (2008).